



## **MEDICATIONS TO AVOID PRIOR TO SURGERY**

Patient instructions prior to surgery

DO NOT TAKE the following medications for two weeks before or after surgery. They promote bleeding. This is not an exhaustive list. If you are in doubt about whether it is safe to stop anyone of the medicines listed below, please contact your local doctor (or the specialist who prescribed the medication) as to whether it is safe to stop the medicine.

ACLIN  
ACTIPROFEN  
AGGRASTAT  
ALEVE  
ANAPROX  
ARTHREXIN  
ASA  
ASASANTIN  
ASPALGIN  
ASPIRIN  
ASPRO  
ASTRIX

BEX  
BRUFEN  
BUFFERIN

CANDYL  
CARDIPRIN  
CARTIA  
CHEMGESIC  
CLINORIL  
CODOX  
CODRAL  
CODRAL PRODUCTS  
CODIPHEN  
CELEBREX

DECRIN  
DISPRIN  
DOLOXENE  
ECOTRIN  
FELDENE

DENAC  
DENSAID

HYCODIN

IBUPROFEN  
INDOCID  
INDOMED  
ISCOVER

MAREVAN  
MEFIC  
MOBILIS  
MORPHALGIN

NAPROGESIC  
NUROFEN  
NUROFEN JUNIOR

ORTHOXICOL  
ORUDIS  
ORUVAIL

PALAPRIN FORTE  
PERCODAN  
PERPAIN  
PERSANTIN  
PIROX  
PLAVIX  
PONSTAN  
PROXEN

RAFEN  
SALDAC  
SOLCODE  
SOLPRIN  
SOLVIN  
SPREN  
S.R.A  
SURGAM

TILCOTIL  
TICLODIPINE  
TILODENE  
TORADIL

VEGANIN  
VINCENTS POWDERS  
VIOXX  
VOLTAREN  
WARFARIN  
WINSPRIN

### **HERBAL AND NATURAL PRODUCTS**

GINGKO  
GARLIC  
ECHINACEA  
OMEGA-3 LIQUID  
FISH OIL  
VITAMIN E  
FEVERFEW  
GINGER  
GINSENG

\* If you have pain or headache it is SAFE to use

CAPADEX  
CODALGIN  
DIFFLAM  
DIGESIC  
DOLOXENE  
DYMADON  
MERSYNDOL

PAINSTOP  
PANADEINE  
PANADEINE FORTE  
PANADOL  
PANAMAX  
PARACETAMOL  
PARADEX

62 Buckley Street  
Essendon 3040  
phone 03 9375 2099  
fax 03 9375 2486  
Provider No 244532VX

Suite 222 / Level 2  
100 Victoria Parade  
East Melbourne 3002  
phone 03 9650 0522  
fax 03 9650 2522  
Provider No 295368BW

ABN 39 450 749081

www.leungent.com.au